



Dizziness Handicap Inventory	YES (4)	SOMETIMES (2)	NO (0)
Because of your problem have you been embarrassed in front of others			
Because of your problem, are you afraid people will think you are intoxicated			
Because of your problem, are you afraid to leave home without company			
Because of your problem, are you afraid to stay home alone			
Because of your problem, are you depressed			
Because of your problem, do you avoid heights			
Because of your problem, do you feel frustrated			
Because of your problem, do you feel handicapped			
Because of your problem, do you have difficulty getting into or out of bed			
Because of your problem, do you have difficulty reading			
Because of your problem, do you restrict travel			
Because of your problem, is it difficult for you to concentrate			
Because of your problem is it difficult for you to go for a walk			
Because of your problem is it difficult for you to walk around the house			
Because of your problem, is it difficult for your to do strenuous work			
Do quick movements of your head increase your problem			
Does bending over increase your problem			
Does looking up increase your problem			
Does performing ambitious activities increase problems			
Does turning over in bed increase your problem			
Does walking down a sidewalk increase your problem			
Does walking down the aisle of a supermarket increase your problems			
Does your problem interfere with job or household responsibilities			
Does your problem significantly restrict your participation in social activities			
Has the problem placed stress on relationships			
TOTAL	TOTAL	/100	

Initially reviewed by Amy M. Yorke, PT, NCS and the MS EDGE task force and Irene Ward, PT, DPT, NCS and the TBI EDGE taskforce of the Neurology Section of the APTA; Updated by Salomi R. Vora in 10/2012. Updated with references for individuals with vestibular disorders by Tracy Rice, PT, MPH, NCS and Jenny Fay, PT, DPT, NCS and the Vestibular EDGE task force of the Neurology Section of the APTA (2013)